

Almost periodic mappings to complex manifolds.

S. Favorov, N. Parfyonova

Abstract. H. Bohr in 1930 proved that if a holomorphic bounded function on a strip is almost periodic on a straight line in the strip, then it is almost periodic on the whole strip. We find some conditions when the result is valid for holomorphic mappings of tube domains to various complex manifolds.

A continuous mapping F of a tube $T_K = \{z = x + iy : x \in \mathbb{R}^m, y \in K \subset \mathbb{R}^m\}$ to a metric space X is *almost periodic* if the family $\{F(z+t)\}_{t \in \mathbb{R}^m}$ of shifts along $\mathbb{R}^m$ is a relatively compact set with respect to the topology of the uniform convergence on T_K .

Further, let X be a complex manifold, and F be a holomorphic mapping of a tube $T_\Omega = \{z = x + iy : x \in \mathbb{R}^m, y \in \Omega\}$, with the convex open base $\Omega \subset \mathbb{R}^m$, to X . We will say that F is *almost periodic* if the restriction of F to each tube T_K , with the compact base $K \subset \Omega$, is almost periodic.

For $X = \mathbb{C}$ we obtain the well-known class of holomorphic almost periodic functions; for $X = \mathbb{C}^m$ the corresponding class was being studied in [3], [7], [8], [6]; for $X = \mathbb{CP}$, we get the class of meromorphic almost periodic functions that was being studied in [4], [10], [5]; the class of holomorphic almost periodic curves, corresponding to the case $X = \mathbb{CP}^m$, was being studied in [6].

The following theorem is well known:

Theorem B (H. Bohr [2]) *If a holomorphic bounded function on a strip is almost periodic on some straight line in this strip, then this function is almost periodic on the whole strip.*

This theorem was extended to holomorphic functions on a tube domain in [11]; besides usual uniform metric, various integral metrics were being studied there.

The direct generalization of Theorem B to complex manifold is not true:

Example. Let $f(z)$ be the quotient of two periodic functions $\sin \sqrt{2}z$ and $\sin z$. It is clear that the restriction of $f(z)$ to any straight line $\{z = x + iy_0 : x \in \mathbb{R}\}$, $y_0 \neq 0$, is almost periodic on this line. Besides, $f(z)$ is bounded as a mapping to the compact manifold $\mathbb{CP}$. Nevertheless, zeros and poles of $f(z)$ are not separated on the real axis $z = x \in \mathbb{R}$, therefore $f(z)$ is not almost periodic on any substrip containing the real axis (see [4]).

In order to give the right version of Theorem B, we need the following proposition.

Proposition. *If K is compact subset of $\mathbb{R}^m$, then an almost periodic mapping F of T_K to a metric space X is uniformly continuous and $F(T_K)$ is a*

relatively compact subset of X .

Proof. Let

$$\varphi(t) = \sup_{y \in K} d(F(z + t), F(z)), \quad t \in \mathbb{R}^m.$$

It is easy to prove that $\varphi(t)$ is an almost periodic function on $\mathbb{R}^m$, therefore the function $\varphi(t)$ extends continuously to Bohr's compactification B of the set $\mathbb{R}^m$ (see, for example, [1]). Hence for any point $\tau \in B$ and any $\varepsilon > 0$ there exists a neighborhood $U \subset B$ of this point such that

$$|\varphi(t) - \varphi(t')| < \varepsilon$$

for each $t, t' \in U \cap \mathbb{R}^m$.

Using the obvious inequality

$$\varphi(t - t') \leq |\varphi(t) - \varphi(t')|$$

we get for all $z \in T_K, t, t' \in U \cap \mathbb{R}^m$

$$d(F(z + t'), F(z + t)) \leq \varepsilon.$$

Therefore the function

$$\overline{F}(\tau, y) = \lim_{x \rightarrow \tau, x \in \mathbb{R}^m} F(x + iy)$$

is well-defined and continuously maps the compact set $B \times K$ to X . Since $\overline{F}(x, y) = F(x + iy)$ for $x \in \mathbb{R}^m$, we obtain all the statements of our proposition.

Note that a bounded holomorphic function on a tube domain is uniformly continuous on every subtube with the compact base, but bounded holomorphic mappings, in general, have no this property. Therefore the following result is natural. **Theorem.** *Let F be a holomorphic mapping of T_Ω to a complex manifold X such that for every compact subset $K \subset \Omega$ the mapping F is uniformly continuous on T_K and $F(T_K)$ is a relatively compact subset of X . If the restriction of $F(z)$ to some hyperplane $\mathbb{R}^m + iy'$ is almost periodic, then $F(z)$ is an almost periodic mapping of T_Ω to X .*

Corollary. *Let F be a holomorphic mapping form T_Ω to a compact complex manifold X such that F is uniformly continuous on T_K for every compact set $K \subset \Omega$. If the restriction of $F(z)$ to some hyperplane $\mathbb{R}^m + iy'$ is almost periodic, then $F(z)$ is an almost periodic mapping of T_Ω to X .* **Proof the Theorem.** Take an arbitrary sequence $\{t_n\} \subset \mathbb{R}^m$. Since the function $F(z)$ is uniformly continuous, the family $\{F(z + t_n)\}$ is equicontinuous on each compact set $S \subset T_\Omega$. Further, it follows from the condition of the Theorem that the union of all the

images of S under mappings of this family is contained in a compact subset of X . Therefore, passing on to a subsequence if necessary, we may assume that the sequence $\{F(z + t_n)\}$ converges to a holomorphic mapping $G(z)$ uniformly on every compact subset of T_Ω . It is easy to see that the mapping $G(z)$ is bounded and uniformly continuous on every tube T_K with the compact base $K \subset \Omega$. Let us prove that this convergence is uniform on every T_K . Assume the contrary. Then we get

$$d(F(z_n + t_n), G(z_n)) \geq \varepsilon_0 > 0 \quad (1)$$

for some sequence $z_n = x_n + iy_n \in T_{K'}$, where K' is some compact subset of Ω . Replacing sequence by a subsequence if necessary, we may assume that the mappings $G(x_n + z)$ converge to a holomorphic mapping $H(z)$, and the mappings $F(z + x_n + t_n)$ converge to a holomorphic mapping $\tilde{H}(z)$ uniformly on every compact subsets of T_Ω . We may also assume that $y_n \rightarrow y_0 \in K'$. Using (1) we get

$$|\tilde{H}(iy_0) - H(iy_0)| \geq \varepsilon_0.$$

Since the mapping $F(x + iy')$ of $\mathbb{R}^m$ to X is almost periodic, we may assume that a subsequence of the mappings $F(x + t_n + iy')$ converges to $G(x + iy')$ uniformly in $x \in \mathbb{R}^m$. Therefore the sequences of mappings $F(x + x_n + t_n + iy')$ and $G(x + x_n + iy')$ have the same limit, i. e., $\tilde{H}(x + iy') = H(x + iy')$ for all $x \in \mathbb{R}^m$. Since $\tilde{H}(z)$, $H(z)$ are holomorphic mappings, we get $\tilde{H}(x + iy) \equiv H(x + iy)$ on T_Ω . This contradiction proves the Theorem.

REFERENCES

- [1] Berg Ch. Introduction to the almost periodic functions of Bohr // Proceedings of a symposium held in Copenhagen, April 24-25. - 1987. - P.15-24.
- [2] Bohr H. Ueber analytische fastperiodische Funktionen // Math. Ann. - 1930 - Vol. 103 -P. 1-14.
- [3] Favorov S.Ju., Rashkovskii A.Ju., Ronkin L.I. Almost periodic divisors in a strip // Journal D'analyse Mathematique. - 1998. - Vol.74 - P.325-345.
- [4] Parfyonova N.D., Favorov S.Yu. Meromorphic almost periodic functions // Math. Stud. Lviv. - 2000. - Vol. 13. - P.190-198.
- [5] Parfyonova N.D. Meromorphic almost periodic functions and their continuous extension on the Bohr's compact // Vestnik Kharkovskogo Universiteta - 2002. - Vol. 542. - P. 73-84. (Russian).

- [6] Parfyonova N.D. Holomorphic almost periodic mappings into complex manifolds // MAG - 2002. - Vol. 9 - No. 2. - P. 294-305. (Russian).
- [7] Rashkovskii A.Yu. Monge-Ampere operators and Jessen functions of holomorphic almost periodic mappings. // MAG - 1998 - No. 5. - P. 274-296.
- [8] Rashkovskii A.Yu. Flows associated with holomorphic almost periodic functions. // MAG - 1995. - Vol. 2, No. 2. - C. 150-169.
- [9] Ronkin L.I. Jessens' theorems for holomorphic almost periodic functions in tube domains. // Sib. Math. Ju. -1987. - Vol. 28, No. 3. - P. 199-204. (Russian).
- [10] Sunyer i Balaguer F. Una nova generalitzacio de les functions gairebe-periodiques. // Inst.d'Estudis Catalans, Arxius de la seccio de ciencies XVII. - 1949 - P.3-46.
- [11] Udodova O.I. Holomorphic almost periodic functions in tube domain // Vestnik Kharkovskogo Universiteta - 2002 - No. 542. - P. 96-105. (Russian).